
Role of Biomarkers and Surrogate Endpoints: Limitations of Serum Concentrations as a Predictor of Response

Shivendra V. Singh

Department of Pharmacology and University of Pittsburgh Cancer Institute

Garlic Derived Sulfur Compounds



Cellular Effects of Garlic Constituents

- DADS-induced apoptosis (HCT116)
 - ↑ intracellular calcium (Milner- 1996)
- DADS-induced G2/M arrest (HCT116)
 - ↓ Cdk1, ↑ p-Cdk1, ↓ Cdc25C, ↑ Cyclin B (Milner-1997)
- DADS-induced apoptosis (MDA-MB-231)
 - ↑ Bax, ↓ Bcl-xL, and caspase-3 activation (Nakagawa- 2001)



Cellular Effects of Garlic Constituents

- DADS-induced apoptosis (SH-SY5Y neuroblastoma)-
 - ↑ ROS, cytochrome c release, JNK activation, and activation of caspase-3 and -9 (Filomeni et al- 2003)

SAMC-induced G2/M arrest and apoptosis induction (SW480, HT-29)-

microtubule depolymerization, JNK and caspase activation (Xiao et al- 2003)

J Natl Cancer Inst 2002 Nov 6;94(21):1648-51

Allium vegetables and risk of prostate cancer: A population-based study.

Hsing AW, Chokkalingam AP, Gao YT, Madigan MP, Deng J, Gridley G, Fraumeni JF Jr. Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD

Epidemiologic and laboratory studies suggest that allium vegetables and garlic constituents have antitumor effects. In a population-based, case-control study conducted in Shanghai, China, we investigated the association between intake of allium vegetables, including garlic, scallions, onions, chives, and leeks, and the risk of prostate cancer. We administered in-person interviews and collected information on 122 food items from 238 case subjects with incident, histologically confirmed prostate cancer and from 471 male population control subjects. Men in the highest of three intake categories of total allium vegetables (>10.0 g/day) had a statistically significantly lower risk (odds ratio [OR] = 0.51, 95% confidence interval [CI] = 0.34 to 0.76; P(trend)<.001) of prostate cancer than those in the lowest category (<2.2 g/day). Similar comparisons between categories showed reductions in risk for men in the highest intake categories for garlic (OR = 0.47, 95%CI = 0.31 to 0.71; P(trend)<.001) and scallions (OR = 0.30, 95% CI = 0.18 to 0.51; P(trend) < .001). The reduced risk of prostate cancer associated with allium vegetables was independent of body size, intake of other foods, and total calorie intake and was more pronounced for men with localized than with advanced prostate cancer.



Structures of Organosulfides

CH₂=CH-CH₂-S-CH₂-CH=CH₂ Diallyl sulfide (DAS)


CH₂=CH-CH₂-S-S-CH₂-CH=CH₂ Diallyl disulfide (DADS)

CH₂=CH-CH₂-S-S-S-CH₂-CH=CH₂ Diallyl trisulfide (DATS)

CH₃-CH₂-CH₂-S-CH₂-CH₂-CH₃ Dipropyl sulfide (DPS)


CH₃-CH₂-CH₂-S-S-CH₂-CH₂-CH₃ Dipropyl disulfide (DPDS)

DATS Inhibits PC-3 Cell Proliferation




DATS Treated PC-3 Cells Are Arrested in G2/M Phase of the Cell Cycle

(24 h exposure)







DATS Treated PC-3 Cells Are Arrested in G2/M Phase of the Cell Cycle











DATS-Induced G2/M Phase Arrest in Synchronized PC-3 Cells






Normal Prostate Epithelial Cells Are Resistant to Cell Cycle Arrest by DATS













DATS Inhibits Cdk1 activity










Nuclear Accumulation of Cyclin B1 in DATS Treated PC-3 Cells






DATS Arrests Cells in Mitosis












Group	Treatment	$\%G_0/G_1$	% S	$\%G_2$ -M
Mock	DMSO	50 ± 5	20 ± 4	25 ± 2
	DATS	12 ± 1*	18 ± 3	58 ± 5*
Control	DMSO	52 ± 4	17 ± 2	25 ± 1
	DATS	17 ± 3*	17 ± 3	53 ± 7*
Chk2	DMSO	53 ± 4	17 ± 3	25 ± 1
	DATS	$18 \pm 1*$	18	$54 \pm 1*$

Data are mean \pm SE (n= 3). Similar results were observed in 2 experiments. *Significantly different compared with control by one-way ANOVA (p < 0.05).




Chk2 is dispensable for DATS-induced Cdc25C and Cdk1 phosphorylation and Mitotic Arrest




mock


control siRNA


Chk2 siRNA






G2/M Arrest in HCT-15 Cells Transfected with WT and Mutant Chk2







DATS Increases Chk1 Phosphorylation




Chk1 Protein Depletion Fails to Overcome DATS-induced G2/M arrest






PI Fluorescence

Chk1 Protein Knockdown Attenuates DATS-Induced Histone H3 Phosphorylation





Chk1 Protein Knockdown Inhibits DATS-Induced Mitotic Arrest




^{*}Significantly different compared with control by one-way ANOVA (P < 0.05)



Conclusions

- DATS causes G2 and M phase cell cycle arrest due to:
 - ROS-dependent degradation of Cdc25C and increased serine-216 phosphorylation of Cdc25C
 - Even though Chk1 protein knockdown inhibits DATS induced phosphorylation of Cdc25C, G₂ arrest is not attenuated.
 - Chk1depletion inhibits DATS-induced M phase arrest (securin, APC?)
- DATS induces apoptosis by:
 - Increasing JNK/ERK mediated phosphorylation of Bcl-2
 - Inhibiting PI3K/Akt pathway



Future Directions

- Mechanism of Chk1-dependent mitotic arrest (securin, Cdc20, APC?).
- Mechanism of DATS induced JNK activation (glutaredoxin/thioredoxin-ASK1?)
- Activity of DATS against prostate tumorigenesis in TRAMP model.
- Toxicological evaluation
- Pharmacokinetics
- Clinical Trial

Thanks to

- CA55589-07, CA76348-06, CA101753-01
- Dr. Dong Xiao
- Dr. Anna Herman-Antosiewicz
- Karen Lew
- Yan Zeng
- Dr. Sanjay K. Srivastava
- Stanley Marynowski
- Dr. Hui Xiao
- Dr. Sung Choi
- Dr. Jedrzej Antosiewicz
- Kamayani Singh